

**SEROXAT/PAXIL
ADOLESCENT DEPRESSION
Position piece on the phase III clinical studies**

EXECUTIVE SUMMARY

Results from the 2 placebo-controlled, phase III clinical trials designed to assess the efficacy and safety of Seroxat/Paxil in adolescents with major depression are now available.

Study 329 (conducted in the US) showed trends in efficacy in favour of Seroxat/Paxil across all indices of depression. However, the study failed to demonstrate a statistically significant difference from placebo on the primary efficacy measures. The second study (study 377), which was conducted in Europe, South America, South Africa and the United Arab Emirates, showed a high placebo response rate and failed demonstrate any separation of Seroxat/Paxil from placebo.

Data from these 2 studies are insufficiently robust to support a label change and will therefore not be submitted to the regulatory authorities. Results from Study 329 will be presented in abstract form at the ECNP meeting (Paris, November 1999) and a full manuscript will be progressed. There are no plans to publish data from Study 377.

SEROXAT/PAXIL
ADOLESCENT DEPRESSION
Position piece on the phase III clinical studies
FOR INTERNAL USE ONLY

SITUATION

2 SB sponsored, placebo-controlled, phase III clinical trials have been conducted, Study 329 (US) and Study 377 (Europe, South America, South Africa and Saudi Arabia), in order to assess the efficacy and safety of Seroxat/Paxil (up to 40mg/day) in the treatment of adolescents (aged between 13 and 18 years and 11 months) with unipolar major depressive disorder (diagnosed according to DSM IIIR, Study 329 or DSM IV criteria, Study 377).

Study 329 was a placebo-controlled, imipramine comparator study with an 8 week acute treatment phase followed by a 6 month extension phase. The acute phase has completed and the extension phase is due to complete at the end of 1998. 275 patients were recruited to the study. Results from the acute phase of this study show that there were no statistically significant differences from placebo on either of the primary efficacy parameters (change from baseline in HAMD total scores and the proportion of responders-where response was defined as a $\geq 50\%$ reduction from baseline in HAMD score or a HAMD score ≤ 8 at endpoint). However, trends in favour of paroxetine compared with placebo were seen across all the indices of depression (change from baseline in HAMD total [p=0.133], HAMD responders [p=0.112], CGI [p=0.094] and K-SADS [p=0.065] scores) and statistically significant differences from placebo were observed in the proportion of patients in remission (defined as a HAMD score of ≤ 8 at endpoint). In general, the response to imipramine was similar to that for placebo. The 6 month extension phase has now completed and is scheduled to report at the end of 1998.

Study 377 was a 12 week placebo-controlled study, conducted in 276 adolescents with major depression. There was a high placebo response rate in this study and no statistically or clinically significant differences from placebo were observed on either of the primary efficacy variables (proportion of patients achieving a $\geq 50\%$ reduction from baseline in total MADRS scores and change from baseline in the K-SADS-L depressive subscale score). The only differences from placebo (secondary efficacy variables) were seen in a subgroup of patients who were ≥ 16 years of age.

Possible explanations for the high placebo response include;

- 1) The large number of study visits
- 2) the duration of the assessments
- 3) The fact that concomitant psychotherapy was not excluded
- 4) Question marks about the adequacy of using currently available diagnostic criteria and rating scales in younger patients
- 5) Adolescents may be more susceptible to a placebo effect
- 6) Developmental issues. Children and adolescents may respond in a pharmacologically different manner due to quantitative and/or qualitative differences in neurotransmitter/receptor systems.

Conclusions from these studies:

- There were no differences in the safety profile of Seroxat/Paxil in adolescents when compared to that already established in the adult population
- The efficacy data from the above clinical trials are insufficiently robust to support a regulatory submission and label change for this patient population.

OTHER DATA:

Ongoing studies: SB France are conducting a locally funded double-blind, comparative study of Seroxat/Paxil with clomipramine in adolescents with major depression (Study 511). In addition, a study in adolescents with OCD (Study 453) is underway in the US. This study comprises a 16 week open label Seroxat/Paxil treatment phase, followed by double-blind, randomisation to paroxetine or placebo for a further 16 weeks of treatment. The regulatory acceptability of these 2 studies needs to be established.

Published data: A review of the literature shows that 2 studies assessing the use of paroxetine in the treatment of 34 adolescents and children with depression have been published (Rey-Sanchez and Gutierrez-Cesares, 1997; Findling et al; 1996).

The first study (Rey-Sanchez and Gutierrez-Cesares, 1997) was a retrospective survey of data from 25 adolescents (aged 13-17 years) treated with paroxetine. Patients were diagnosed according to ICD 10 criteria. In 13 of the patients unipolar major depression was not the primary diagnosis. 17 patients received paroxetine as a monotherapy, 8 also received concomitant psychotropic medications (n=7 benzodiazepines, n=1 haloperidol). Paroxetine was administered at doses of 10mg (14 patients) or 20mg/day (11 patients). No specific depression rating scales were used, response was based on clinical judgement. 76% patients

had a satisfactory response (11 complete remission, 8 improved with residual symptoms). A lack of satisfactory response in was observed in 6 (24%) patients. Eight patients reported side effects (somnolence or sleep disorders n=6, asthenia n=4, nausea n=3, tachycardia n=2, diarrhea n=2, headache n=2, orthostatic hypotension n=1; restlessness n=1). Two patients were withdrawn due to one due to anxiety, one due to hypotension and dizziness)

The second study (Findling et al; 1996) was conducted in 9 patients aged between 7-15 years (children and adolescents) meeting DSM IV criteria for a major depressive disorder. Symptomatology was assessed using HAM-D for subjects aged 13 to 15 years, and the childhood depression rating scale (CDRS) subjects aged 12 or younger. Paroxetine was initially given at a dose of 10mg/day. This was escalated to 20mg/day if the patient had not responded after 4 weeks of treatment. 8/9 patients responded to treatment with paroxetine. Three patients had complete remission, 5 patients had a >50% reduction in total CDRS score from baseline. CGI improved in all patients. One patient withdrew from the study at week 2 due to an adverse experience. This patient was found to have elevated serum paroxetine levels and was a poor 2D6 metaboliser. Assessment of pharmacokinetic parameters in this study showed that paroxetine had a similar half life to that reported in the adult population (15.7h [sd 9.0h] vs 24h, respectively).

COMPETITOR ACTIVITIES:

Lilly are believed to be in near to completing their phase III clinical trials in adolescent depression. One relatively large placebo-controlled 8 week study with an open 12 month follow-up period conducted in 96 patients (aged 8-18 years) has recently been published (Emslie et al; 1997 and 1998). These data show that 56% (27/48) patients on fluoxetine (20mg/day) compared with 33% (16/48) patients on placebo were rated as much or very much improved on the CGI at Week 6 (p=0.02). In the 12 month follow-up period, 85% (n=74) patients recovered from the depressive episode (47 on fluoxetine, 22 on placebo and 5 on other antidepressants or lithium). Twenty nine (39%) of the patients (36% of those who had recovered on fluoxetine [17/47] and 41% of those who had recovered on placebo [9/22] had a recurrence of depression during the 12 month follow-up (a higher recurrence rate than seen in adults). Other published data on fluoxetine are from small open studies or individual case reports (Colle et al; 1994).

Pfizer already have positive data (including PK data) and are licenced in the US for the treatment of adolescent OCD. In addition, Pfizer are also believed to be conducting clinical trials in adolescent depression. Available published data are

limited, derived from small open studies in adolescent depression (McConville et al; 1996; Tierney et al; 1995)

TARGET

To effectively manage the dissemination of these data in order to minimise any potential negative commercial impact.

PROPOSALS

- Based on the current data from Studies 377 and 329, and following consultation with SB country regulatory and marketing groups, no regulatory submissions will be made to obtain either efficacy or safety statements relating to adolescent depression at this time. However data (especially safety data) from these studies may be included in any future regulatory submissions, provided that we are able to go on and generate robust, approvable efficacy data. The rationale for not attempting to obtain a safety statement at this time is as follows;
 - i) regulatory agencies would not approve a statement indicating that there are no safety issues in adolescents, as this could be seen as promoting off-label use
 - ii) it would be commercially unacceptable to include a statement that efficacy had not been demonstrated, as this would undermine the profile of paroxetine.
- Positive data from Study 329 will be published in abstract form at the ECNP (Paris, November 1998) and a full manuscript of the 329 data will be progressed.
- The regulatory acceptability of Studies 511 and 453 and any other data in this patient population will continue to be investigated.

REFERENCES

1. Rey-Sanchez F; Gutierrez-Cesares JR. Paroxetine in children with major depressive disorder: an open trial. *J Am Acad Child Adolesc Psychiatry* 1997; 36 (10): 1443-1447.
2. Findling R Fiala S, Myers C et al. Putative determinants of paroxetine response in pediatric patients with major depression. *Psychopharmacol Bull*; 1996; 32 (3):446.
3. Emslie GJ, Rush AJ, Weinberg WA et al. A double-blind, randomized, placebo-controlled trial of fluoxetine in children and adolescents with depression. *Arch Gen Psychiatry*; 1997; 54 (11): 1031-1037.
4. Emslie GJ, Rush AJ, Weinberg WA et al. Fluoxetine in child and adolescent depression: acute and maintenance treatment. *Depression and Anxiety (US)*; 1998; 7 (1): 32-39.
5. Colle LM, Belair JF, DiFeo M et al. Extended open label fluoxetine treatment of adolescents with major depression. *J Child Adolesc Psychopharmacol*. 1994; 4 (4): 225-232.
6. McConville BJ, Minnery RL, Sorter MT et al. An open study of the effects of sertraline on adolescent major depression. *J Child Adolesc Psychopharmacol*. 1996; 6 (1):41-51
7. Tierney E, Joshi PT, Llinas JF et al. Sertraline for major depression in children and adolescents: Preliminary clinical experience. *J Child Adolesc Psychopharmacol*; 1995; 5 (1):13-27.